

Enzymatic activity with an incomplete catalytic spine: insights from a comparative structural analysis of human CK2 α and its paralogous isoform CK2 α'

Nils Bischoff · Jennifer Raaf · Birgitte Olsen ·
Maria Bretner · Olaf-Georg Issinger ·
Karsten Niefind

Received: 13 June 2011 / Accepted: 24 June 2011
© Springer Science+Business Media, LLC. 2011

Abstract Eukaryotic protein kinases are fundamental factors for cellular regulation and therefore subject of strict control mechanisms. For full activity a kinase molecule must be penetrated by two stacks of hydrophobic residues, the regulatory and the catalytic spine that are normally well conserved among active protein kinases. We apply this novel spine concept here on CK2 α , the catalytic subunit of protein kinase CK2. *Homo sapiens* disposes of two paralog isoforms of CK2 α (*hsCK2 α* and *hsCK2 α'*). We describe two new structures of *hsCK2 α* constructs one of which in complex with the ATP-analog adenylyl imidodiphosphate and the other with the ATP-competitive inhibitor 3-(4,5,6,7-tetrabromo-1*H*-benzotriazol-1-yl)propan-1-ol. The former is the first *hsCK2 α* structure with a well defined cosubstrate/magnesium complex and the second with an open β 4/ β 5-loop. Comparisons of these structures with existing CK2 α /CK2 α' and cAMP-dependent protein kinase (PKA) structures reveal: in *hsCK2 α'* an open conformation of the interdomain hinge/helix α D region that is critical for ATP-binding is found corresponding to an incomplete catalytic spine. In contrast *hsCK2 α* often adopts the canonical, PKA-like version of the catalytic spine which

correlates with a closed conformation of the hinge region. *HsCK2 α* can switch to the incomplete, non-canonical, *hsCK2 α'* -like state of the catalytic spine, but this transition apparently depends on binding of either ATP or of the regulatory subunit CK2 β . Thus, ATP looks like an activator of *hsCK2 α* rather than a pure cosubstrate.

Keywords Protein kinase CK2 · Casein kinase 2 · Eukaryotic protein kinases · Regulatory spine · Catalytic spine · Paralog isoforms of human CK2 α

Introduction

Eukaryotic protein kinases (EPKs) are central regulators of cellular key processes and require for this function a strict control of their own catalytic activity. Consequently the elucidation of regulatory mechanisms of EPKs and their structural bases are fundamental subjects of EPK research. The growing knowledge about these mechanisms and their common structural principles are reflected in a series of comprehensive reviews [1–4].

The latest approach to gain a thorough understanding of the structural constraints that govern the active state of EPKs was introduced by Kornev et al. [5] on the basis of “Local spatial pattern alignment”, a novel graph theoretical method for structural comparison [6]. According to this concept a fully active EPK requires two stacks of hydrophobic residues—called regulatory R-spine and catalytic C-spine—that start from the hydrophobic helix α F in the centre of the C-terminal EPK domain, penetrate the catalytic core and thereby cross the interface between the two main domains (Fig. 1). A continuous R-spine is the fundamental structural condition for catalytic activity of an EPK whereas the C-spine can be slightly disrupted by

Nils Bischoff and Jennifer Raaf—Common first authors.

N. Bischoff · J. Raaf · K. Niefind (✉)
Universität zu Köln, Institut für Biochemie, Zülpicher Straße 47,
50674 Köln, Germany
e-mail: Karsten.Niefind@uni-koeln.de

B. Olsen · O.-G. Issinger
Syddansk Universitet, Institut for Biokemi og Molekylær
Biologi, Campusvej 55, 5230 Odense, Denmark

M. Bretner
Faculty of Chemistry, Warsaw University of Technology,
Noakowskiego 3, Warsaw, Poland




Fig. 1 Catalytic spine (C-spine; yellow) and regulatory spine (R-spine; red) of cAMP-dependent protein kinase (PKA). The “gatekeeper” is a methionine residue in PKA (Met120), but Phe113 in *hsCK2 α* . Reprinted from *Biochim Biophys Acta*, Vol. 1804, Kornev AP and Taylor SS, “Defining the conserved internal architecture of a protein kinase”, pp 440–444, 2010, with kind permission from Elsevier. (Color figure online)

domain motions in the course of the catalytic cycle and is fully established only after integration of the adenine base of the cosubstrate ATP.

In the context of protein kinase CK2—an ubiquitous, essential and pleiotropic Ser/Thr kinase composed of two catalytic subunits (CK2 α) attached to a dimer of non-catalytic subunits (CK2 β)—the spine concept was applied for the first time by Prudent et al. [7]. These authors hypothesized that the effect of polyoxometalates—a group of highly effective non-ATP-competitive CK2 inhibitors—might consist in a disruption of the R-spine of CK2 α which is normally well conserved (Fig. 2c). We ourselves used the R-spine concept recently in the context of a structural study on human CK2 α' [8]. There we interpreted a hydrophobic pocket at the surface of the N-terminal domain that can be filled with a conserved tryptophan residue (Trp34 in CK2 α') and that is equivalent to the PIF-pocket of PKA and other AGC family kinases [9] as an extension of the R-spine.

Here we will present a closer look at the C-spine of the catalytic CK2 subunit. CK2 has been called “a challenge to the canons” [10], a qualification that is also valid with respect to the C-spine. This is indirectly apparent from Fig. 2a/b which stems from a recent review by Taylor and Kornev [4]: in Fig. 2a the conservation of the R-spine is demonstrated by an overlay of 23 active EPK structures (plus 2 structures of prokaryotic EPK-like kinases) among them CK2. In Fig. 2b, however, in which Taylor and Kornev [4] illustrated the canonical C-spine, CK2 was left out. Obviously CK2 does not fit to the C-spine canon, and we will show here why.

Materials and methods

Crystal structures

For the C-spine analysis we used several crystal structures mainly of human CK2 α (*hsCK2 α*), but also of human CK2 α' (*hsCK2 α'*), maize CK2 α , human CK2 holoenzyme and cAMP-dependent protein kinase (PKA). *hsCK2 α* and *hsCK2 α'* are the two paralogous isoforms of CK2 α encoded in the human genome [11]. Their amino acid sequences are to 90.7% identical up to position 330 whereas the C-terminal segments are completely unrelated (Fig. 3).

The identities of the compared structures can be seen in Fig. 2e. Their Protein Data Bank codes and references are as follows: 3NSZ [12], 2PVR [13], 3OFM [8], 1LP4 [14], 1JWH [15], 3JUH [16], 3BQC [17], 3H30 [18], 3FWQ [19] and 1ATP [20]. Two further structures included in the comparison had not been published previously; the details of their determination are described below.

The first of these structure, a co-crystal structure of *hsCK2 α* ^{1–335} with the ATP-competitive inhibitor 3-(4,5,6,7-tetrabromo-1*H*-benzotriazol-1-yl)propan-1-ol (abbreviated here as “MB002”) [21] that was recently crystallized together with the *hsCK2 α'* mutant *hsCK2 α'* ^{Cys336Ser} [8], was determined to allow a structural comparison of both CK2 α paralogs with the same ligand at the ATP site. The second novel structure contains a chimeric construct comprising residues 1–325 of *hsCK2 α* and residues 327–350 of *hsCK2 α'* (see residues in bordered boxes of Fig. 3) and is called *hsCK2 α α'* from here on. We explained the rationale behind *hsCK2 α α'* elsewhere [8]; briefly, the motivation was to determine the structure of the C-terminal region of *hsCK2 α'* in which it is unrelated to *hsCK2 α* (Fig. 3).

Mutagenesis

The construction of the C-terminally truncated mutant *hsCK2 α* ^{1–335} was described by Ermakova et al. [22]. For the construction of *hsCK2 α α'* the DNA sequence coding for the kinase core of *hsCK2 α* (amino acids 1–325) was amplified with oligonucleotides introducing a *NdeI* restriction site (forward primer: 5'-GGAATTCCATATGT CGGGACCCGTGCCAAG-3') and *SnaBI* plus *HindIII* restriction sites (reverse primer: 5'-CCCAAGCTTTAC GTAGAAATAGGGGTGCTCCATTGCC-3'). The fragment was ligated into the *NdeI* and *HindIII* cut pT7-7 expression vector.

To obtain the C-terminus of *hsCK2 α'* , complementary oligonucleotides with a 3'-end overlap suitable for ligation with a *HindIII*-cut restriction site were used (P-5'-CCT GTGGTGAAGGAGCAGTCCCAGCCTTGTGCAGACA ATGCTGTGCTTTCCAGTGGTCTCACGGCAGCAGC ATG-3' and P-5'-AGCTTTCATCGTGCTGCCGTGAG




Fig. 2 Multiple alignments of residues forming the catalytic spine and the regulatory spine. Illustration of the R-spine (a) and the C-spine (b) as published by Taylor and Kornev [4]. The residue numbers refer to cAMP-dependent protein kinase (PKA). This part of the figure was reprinted from Trends Biochem Sci, Vol. 36, Taylor SS and Kornev AP, “Protein kinases: evolution of dynamic regulatory proteins”, pp 65–77, 2011, with kind permission from Elsevier. R-spine (c) and C-spine (d) in CK2 α . Only in the closed hinge/helix

α D conformation Phe121 overlaps with Met128 of PKA. In part (d) the CK2 β interaction “hot spots” Leu41 and Phe54 [36] were drawn in addition to the C-spine residues (grey background). The colour code of the superimposed structures is given in (e). The PKA structure 1ATP [20] was added as a reference (red side chains and labels). The CK2 α residues were numbered according to the *hsCK2 α* sequence. (Color figure online)




Fig. 3 Sequence alignment of C-terminal regions of CK2 α subunits. The C-terminal segments of human CK2 α (*hsCK2 α*) and CK2 α' (*hsCK2 α'*) differ completely in length and sequence. Both paralogs are significantly longer than maize CK2 α for which the majority of CK2 α crystal structures exist [32]. The bordered boxes indicate the

sequence of the *hsCK2 α'* construct the structure of which was described here (Table 1). The highlighted SP- and TP-motifs in the *hsCK2 α* sequence were reported to be phosphorylated during mitosis [37]

ACCACTGGAAAGCACAGCATTGTCTGCACAAGG CTGGGACTGCTCCTTACCACA-3'). The annealing reaction mixture contained both oligonucleotides and 100 mM potassium acetate, 30 mM HEPES pH 7.4 and 2 mM magnesium acetate and was incubated at 95°C for 4 min followed by an incubation at 70°C for 10 min. After that the reaction mixture was slowly cooled down to room temperature to ensure annealing of the complementary oligonucleotides. The annealed fragment was then ligated into the *SnaBI/HindIII* cut pT7-7-*hsCK2 α ¹⁻³²⁵*-plasmid adjacent to the *hsCK2 α ¹⁻³²⁵*-sequence.

Enzyme production and crystallization

The construct *hsCK2 α'* and the established mutant *hsCK2 α ¹⁻³³⁵* [22] were prepared according to a protocol described previously [13]. For *hsCK2 α'* an additional gel-filtration run was necessary. The purified proteins were concentrated and rebuffed in 500 mM NaCl and 25 mM Tris/HCl, pH 8.5, by ultrafiltration using AMICON Ultra-15 tubes.

Crystallization experiments were performed at 20°C with the sitting drop variant of the vapour diffusion technique (sitting drop). For *hsCK2 α'* initial crystallization experiments with “Crystal Screen Cryo” from Hampton Research led to first crystals. The crystallization conditions were optimized by varying the concentrations of the reservoir components and using “Detergent Screen 2” from Hampton Research. The composition of the optimized crystallization drop was as follows: 0.8 μ l *hsCK2 α'* (12.6 mg/ml), 1.5 μ l 5 mM AMPPNP, 1.5 μ l 10 mM MgCl₂, 1.5 μ l 1 mM CK2 substrate peptide (sequence RRRADDSDDDDD), 0.5 μ l 10% (v/v) Anapoe[®]305, 0.8 μ l reservoir solution (0.17 M ammonium sulfate, 0.1 M sodium cacodylate, pH 6.2, 15% PEG 8000 and 15%

glycerol). Cryo conditions were obtained by reequilibration after changing the reservoir to 0.17 M ammonium sulfate, 0.1 M sodium cacodylate, pH 6.5, 25% PEG 8000 and 25% glycerol.

HsCK2 α ¹⁻³³⁵ was co-crystallized with 3-(4,5,6,7-tet-rabromo-1*H*-benzotriazol-1-yl)propan-1-ol (MB002) [21], i.e., with the same ATP-competitive inhibitor as used before in a co-crystal structure determination with *hsCK2 α ^{Cys336Ser}* [8]. Prior to crystallization 99 volume parts of a protein stock solution that contained 6 mg/ml *hsCK2 α ¹⁻³³⁵* in 0.5 M NaCl, 25 mM Tris/HCl, pH 8.5 was mixed with 1 volume part 10 mM MB002 in dimethyl sulfoxide. After 30 min incubation at room temperature the *hsCK2 α ¹⁻³³⁵*/MB002 mixture was used for vapour diffusion crystallization (sitting drop). In the most successful setup the reservoir contained 4 M NaCl, 0.1 M sodium citrate buffer, pH 5.0. In the drop 0.5 μ l of this reservoir solution was mixed with 0.5 μ l *hsCK2 α ¹⁻³³⁵*/MB002 mixture. *HsCK2 α ¹⁻³³⁵*/MB002 crystals grown under these conditions could be directly frozen in liquid nitrogen and mounted for X-ray diffraction.

X-ray diffraction data collection and structure determination

In both cases X-ray diffraction data were collected at a wavelength of 0.91841 Å and a temperature of 100 K at beamline BL-1 at the BESSY synchrotron in Berlin (Germany). The diffraction data were processed with XDS [23]. The structures were solved by molecular replacement using PHASER [24] and refined with REFMAC [25] from the CCP4 program suite [26]. A parameter file for the MB002 ligand [21] was calculated with PRODRG [27]. Manual model building was performed with COOT [28].

Results and discussion

Quality and overview of the two novel structures

The two novel co-crystal structures (*hsCK2 α ^{1–335}* plus MB002 and *hsCK2 α α'* plus AMPPNP) were refined to reasonable R-factors and stereochemical geometries (Table 1). In spite of the low resolution the *hsCK2 α α'* structure fits well to its final electron density; however, the original aim—to see an ordered structure in the C-terminal region originating from *hsCK2 α'* —was not reached due to disorder after Glu330. Hence, the C-terminal segments of

Table 1 Overview of X-ray diffraction data and refinement

	<i>hsCK2$\alpha$$\alpha'$</i> / AMPPNP	<i>hsCK2α^{1–335}</i> / MB002
Collection and statistics		
Data collection temperature (K)	100	100
Wavelength (Å)	0.91841	0.91841
Space group	P4 ₃	P4 ₃
Lattice constants		
a, b, c, (Å)	127.79, 127.79, 125.31	72.01, 72.01, 133.12
α , β , γ (°)	90.0, 90.0, 90.0	90.0, 90.0, 90.0
Resolution: complete (highest shell) (Å)	38.50–3.00 (3.08–3.00) ¹	19.75–2.30 (2.36–2.30) ¹
R _{sym} (%)	12.7 (85.7) ^a	6.5 (83.0) ^a
Signal to noise ratio (I/ σ ₁)	13.35 (2.91) ^a	20.41 (2.47) ^a
No. of unique reflections	40400 (2957) ^a	30062 (2172) ^a
Completeness (%)	99.9 (100.0) ^a	99.7 (98.4) ^a
Redundancy	7.5 (7.7) ¹	7.6 (7.3) ^a
B-factor from Wilson plot (Å ²)	59.6	54.3
Refinement		
Resolution (Å)	38.50–3.00	19.75–2.30
No. of reflections in working set/test set	39113/1259	29111/912
R _{work} /R _{free}	16.3/20.3	17.0/22.8
No. of atoms/average B-factors (Å ²)		
Protein	11120/61.8	5634/52.6
Ligand/ion	107/53.5	31/73.7
Water	32/42.0	145/41.5
RMS deviations		
Bond lengths (Å)	0.006	0.006
Bond angles (°)	0.915	0.924
Ramachandran plot quality ² /No. of residues in		
Most favoured regions	1027 (86.2%)	530 (87.2%)
Additionally allowed regions	157 (13.2%)	72 (11.8%)
Generously allowed regions	4 (0.3%)	5 (0.8%)
Disallowed regions	4 (0.3%)	1 (0.2%)

^a The values in brackets refer to the highest resolution shell

both *hsCK2 α* and *hsCK2 α'* remain structurally uncharacterized similar to all previous studies in which full-length constructs were used [8, 15, 29]. Probably it requires specific interaction partners to induce ordered conformations in these regions.

Nevertheless the *hsCK2 α α'* structure (basically thus a C-terminally truncated *hsCK2 α* structure) turned out to be valuable for this analysis since in two of the four subunits within its asymmetric unit it contains—as the first human CK2 α structure—a complete and well ordered AMPPNP molecule plus two magnesium ions (Fig. 4a). The whole arrangement is fully functional and similar to the ternary complex of PKA, an inhibitor peptide and ATP (black bonds in Fig. 4a) [20], meaning the γ -phospho group is correctly placed for the transfer reaction. In the two other subunits no ligands are visible so that they can be regarded as apo-forms.


In all four chains of the *hsCK2 α α'* structure the β 4/ β 5-loop as a central element of the CK2 β interface adopts an open and stretched conformation which is present also in the CK2 holoenzyme [15]. Since hitherto, in all structures of unbound (CK2 β -free) *hsCK2 α* the β 4/ β 5-loop was found with a closed conformation, incompatible with CK2 β binding, this is a remarkable detail. It is, however, not really surprising since in the recently published near-atomic resolution structure of *hsCK2 α ^{2–335}* in complex with an ADP-analog the β 4/ β 5-loop was also in the open state [12].

In the case of the *hsCK2 α ^{1–335}*/MB002 complex structure only the ATP site of chain A is filled by the inhibitor while chain B is the second subunit within the asymmetric unit shows an uncomplexed apo-state. In this structure the β 4/ β 5-loop conformation differs between the two chains, i.e., it is open in one subunit and closed in the other which shows that the crystalline environment is a major determinant of the loop conformation.

Differences between *hsCK2 α* and *hsCK2 α'* in MB002 binding

MB002 is so far the only ATP-competitive CK2 inhibitor which has been crystallized with both paralogous forms of human CK2 α . A superimposition of these structures (Fig. 4b) reveals that the binding mode of MB002 differs in both cases: in complex with *hsCK2 α' ^{Cys336Ser}* [8] (black carbon atoms in Fig. 4b) the propanol tail of the inhibitor points to the inner region of the ATP site and forms a hydrogen bond to the conserved water molecule pointed out by Battistutta et al. [30]; in contrast when MB002 is bound to *hsCK2 α ^{1–335}* (yellow carbon atoms in Fig. 4b) the propanol moiety approaches the glycine-rich loop (ATP-binding loop; not shown in Fig. 4b). Here, the conserved lysine Lys68 bends significantly in the direction of

Fig. 4 Stereo pictures to illustrate conformations and ligands at and around the ATP-binding site of *hsCK2 α* and *hsCK2 α'* . In all panels of the figure the illustrated part of the respective main structure (yellow carbon atoms) was embedded in the final electron density with cutoff level 1 σ . The figure was prepared with BOBSCRIPT [38] in combination with Raster3D [39]. (a) Adenylyl imidodiphosphate (AMPPNP) plus Mg^{2+} ions bound to the ATP site of the chimeric construct *hsCK2 α α'* . From the enzyme the catalytic loop and three C-spine residues (Met163, Val66, Val53) were drawn. For comparison ATP plus Mn^{2+} ions from the PKA ternary complex structure 1ATP [20] were added after structural superimposition of the protein matrices. (b) The ATP-competitive inhibitor 3-(4,5,6,7-tetrabromo-1H-benzotriazol-1-yl)propan-1-ol (MB002) [21] bound either to *hsCK2 α $^{1-335}$* (yellow carbon atoms; thick bonds) or to *hsCK2 α' Cys336Ser* (PDB file 3OFM; black carbon atoms; thin bonds) [8]. (c) Structural ambiguity of Phe121 and Tyr125 in *hsCK2 α* . The “closed hinge/canonical C-spine”-state was drawn with black carbon atoms and is represented by the *hsCK2 α $^{1-335}$* /MB002 structure (Table 1; Fig. 4b). The alternative “open hinge/non-canonical spine”-conformation (yellow carbon atoms) was extracted from the *hsCK2 α α'* /AMPPNP structure (Table 1; Fig. 4a). (Color figure online)



the inhibitor and forms a halogen bond to one of the bromine substituents.

A further halogen bond to the MB002 inhibitor not found in the *hsCK2 α ^{Cys336Ser}/MB002* complex is established by the carbonyl oxygen of Asn117 (Fig. 4b). The region around Asn117 is part of the interdomain hinge/helix α D region for which two principle conformations—an open and a closed one—are known [18]. “Open” means that relatively much space is left in the ribose binding region, a necessary condition for the typical “dual cosubstrate specificity” of CK2 [14]. The open conformation was found both in the CK2 holoenzyme [15] and in two maize CK2 α structures with ATP- and GTP-analogs [14] showing a similarly well defined state of the nucleotides and adjacent magnesium ions as seen in Fig. 4a. These findings suggest a correlation between the open hinge/helix α D region conformation and a fully functional way of ATP or GTP binding whereas the closed conformation seems to be associated with a non-productive ATP-binding mode [16, 31]. In Fig. 4b the open state is represented by the *hsCK2 α ^{Cys336Ser}/MB002* complex [8] (black carbon atoms in Fig. 4b) and the closed one by the novel *hsCK2 α ^{1–335}/MB002* complex (Table 1).

The structural ambiguity in the protein environment of the MB002 inhibitor shows that the latter is not able to enforce a particular protein conformation. In fact, as we pointed out recently [8] *hsCK2 α '* just like maize CK2 α prefers the open hinge conformation irrespective of the ligand, the crystallization condition and the crystalline environment. In contrast *hsCK2 α* is so far the only CK2 α protein in which either of the two conformations have been found [32]. Significantly the two novel structures presented here (Table 1) reflect this conformational plasticity: one of them, *hsCK2 α ^{1–335}/MB002* (Fig. 4b), belongs to the cluster of closed hinge structures, but the other one, *hsCK2 α '/AMPPNP* (Fig. 4a), to the open ones.

Phe121: a switchable member of the catalytic spine

The most conspicuous conformational difference in the protein matrices surrounding the two MB002 ligands refers to Phe121 (Phe122 in *hsCK2 α '*) (Fig. 4b). The ambivalent character of this residue in *hsCK2 α* , its ability to switch from a buried position in the closed state to a more solvent exposed one in the open conformation and the correlation of this switch to the productivity of ATP-binding have been noted and discussed before [18, 31, 32]. Now, the novel spine paradigm [4] provokes an interpretation in the light of this concept since on a sequence level [33] Phe121 is equivalent to Met128 of PKA which is a member of the catalytic spine (Figs. 1, 2b–d).

As illustrated by Fig. 1 PKA-Met128 is located at the beginning of the small surface helix α D. Its side chain

forms the centre of a hydrophobic cluster whose further components are the other C-spine members Leu227, Met231, Leu172 and Ile174 (Fig. 1, 2b). Figure 2b emphasizes the structural conservation of this hydrophobic ensemble.

The more surprising is the fact that Kornev and Taylor [4] left CK2 α out from Fig. 2b although including it in Fig. 2a. The reason is indicated in Fig. 2d and more precisely in Fig. 4c: CK2 α fits only with the closed hinge/helix α D conformation which does not correspond to the functional arrangement in Fig. 4a to the C-spine canon; for full functionality, however, it requires the open hinge/helix α D state in which the C-spine is incomplete in the sense that the central (Met128-equivalent) position of the hydrophobic cluster remains more or less unoccupied. “More or less” means that in this case Tyr125 partially fills the relevant space (Fig. 4c). That is why Prudent et al. [8] incorporated Tyr125 in a picture illustrating the two spines of CK2 α . In fact, the “local spatial pattern alignment” procedure [6] underlying the spine concept does not take into account sequence or structural homology so that it is principally justified to include a non-equivalent residue. However, we prefer to classify the C-spine corresponding to the open hinge conformation as “incomplete” since Tyr125 is less hydrophobic than Phe121 and since it occupies only to a minor extend the PKA-Met128 equivalent space at the centre of the C-spine hydrophobic cluster and even this only with structural strain, i.e., with an unfavourable side chain conformation.

Correlations and conclusions

According to the spine concept ATP is not a pure cosubstrate of the kinase reaction. Rather it is, as recently shown for PKA [34], additionally an activating factor since a correctly positioned adenine moiety is required to connect the N-lobe and the C-lobe parts of the catalytic spine (Fig. 1). Against this background it is remarkable that the incomplete and non-canonical C-spine corresponding to an open hinge/helix α D conformation was found in *hsCK2 α* whenever co-crystallized with an ATP- or ADP analog (Fig. 2d, e). In particular the most recent *hsCK2 α* structures, the above-mentioned high-resolution structure 3NSZ [12] and the *hsCK2 α '*-structure of this work, suggest that ATP, ADP or its analogs stabilize the incomplete C-spine state or at least select it for binding. The only apparent counter-example is the complex structure of the mutant *hsCK2 α ^{1–335,Val66Ala,Met163Leu}* with AMPPNP [16]. Here, specific mutations render the ATP site and the C-spine “PKA-like” which supports the combination of canonical C-spine (Fig. 2d, e) and closed hinge conformation. This shift of a conformational preference correlates with a partial breakdown of the dual-cosubstrate specificity, i.e. ATP

is now clearly favoured as a cosubstrate compared to GTP [16].

Taken together, in CK2 α molecules the R-spine is absolutely conserved (Fig. 2c), but the C-spine can be ambiguous (Fig. 2d). For full functionality CK2 α requires an incomplete and non-canonical catalytic spine which is necessarily connected with an open conformation of the hinge/helix α D region. This special structural arrangement seems to be an inherent feature of the maize CK2 α molecule, since in more than 40 crystal structures it was never observed with the alternative, i.e., with a closed hinge region combined to a canonical C-spine. The same may be true for *hsCK2 α'* but given the limited set of two structures [8, 35] it is too early for a generalization.

In the case of its paralog *hsCK2 α* , however, specific interaction partners are required to stabilize the non-canonical C-spine state. It was mentioned already [31] that CK2 β is probably such a factor, and it is noteworthy that the two recently identified CK2 β interaction “hot spot” residues Leu41 and Phe54 [36] are located in direct prolongation of the C-spine axis across the plane of the N-lobe β -sheet (Fig. 2b). The analysis presented here suggests that ATP after CK2 β might be the second stabilizing and activating interaction partner of *hsCK2 α* .

Protein data bank accession codes

The two structures of Table 1 are available from the Protein Data Base (www.rcsb.org) under the accession codes 3RP0 (*hsCK2 α _ α' /AMPPNP*) and 3RPS (*hsCK2 α ^{1–335}/MB002*).

Acknowledgments We are grateful to the staff of the BESSY synchrotron in Berlin, Germany, for assistance with X-ray diffraction experiments. The work was funded by the “Deutsche Forschungsgemeinschaft” (Grant NI 643/4-1) and by the “Forskningsråd for Natur og Univers” (Grant 272-07-0257).

References

- Johnson LN, Noble ME, Owen DJ (1996) Active and inactive protein kinases: structural basis for regulation. *Cell* 85:149–158
- Huse M, Kuriyan J (2002) The conformational plasticity of protein kinases. *Cell* 109:275–282
- Nolen B, Taylor S, Ghosh G (2004) Regulation of protein kinases: controlling activity through activation segment conformation. *Mol Cell* 15:661–675
- Taylor SS, Kornev AP (2011) Protein kinases: evolution of dynamic regulatory proteins. *Trends Biochem Sci* 36:65–77
- Kornev AP, Taylor SS, Ten Eyck LF (2008) A helix scaffold for the assembly of active protein kinases. *Proc Natl Acad Sci USA* 105:14377–14382
- Kornev AP, Taylor SS, Ten Eyck LF (2008) A generalized allosteric mechanism for cis-regulated cyclic nucleotide binding domains. *PLoS Comput Biol* 4:e1000056
- Prudent R, Sautel CF, Cochet C (2010) Structure-based discovery of small molecules targeting different surfaces of protein-kinase CK2. *Biochim Biophys Acta* 1804:493–498
- Bischoff N, Olsen B, Raaf J, Bretner M, Issinger OG, Niefind K (2011) Structure of the human protein kinase CK2 catalytic subunit CK2 α' and interaction thermodynamics with the regulatory subunit CK2 β . *J Mol Biol* 407:1–12
- Biondi RM, Cheung PC, Casamayor A, Deak M, Currie RA, Alessi DR (2000) Identification of a pocket in the PDK1 kinase domain that interacts with PIF and the C-terminal residues of PKA. *EMBO J* 19:979–988
- Pinna LA (2002) Protein kinase CK2: a challenge to canons. *J Cell Sci* 115:3873–3878
- Pyerin W, Ackermann K (2003) The genes encoding human protein kinase CK2 and their functional links. *Prog Nucleic Acid Res Mol Biol* 74:239–273
- Ferguson AD, Sheth PR, Basso AD, Paliwal S, Gray K, Fischmann TO, Le HV (2011) Structural basis of CX-4945 binding to human protein kinase CK2. *FEBS Lett* 585:104–110
- Niefind K, Yde CW, Ermakova I, Issinger OG (2007) Evolved to be active: sulfate ions define substrate recognition sites of CK2 α and emphasise its exceptional role within the CMGC family of eukaryotic protein kinases. *J Mol Biol* 370:427–438
- Niefind K, Pütter M, Guerra B, Issinger O-G, Schomburg D (1999) GTP plus water mimic ATP in the active site of protein kinase CK2. *Nat Struct Biol* 6:1100–1103
- Niefind K, Guerra B, Ermakova I, Issinger O-G (2001) Crystal structure of human protein kinase CK2: insights into basic properties of the CK2 holoenzyme. *EMBO J* 20:5320–5331
- Yde CW, Ermakova I, Issinger O-G, Niefind K (2005) Inclining the purine base binding plane in protein kinase CK2 by exchanging the flanking side-chains generates a preference for ATP as a cosubstrate. *J Mol Biol* 347:399–414
- Raaf J, Klopffleisch K, Issinger O-G, Niefind K (2008) The catalytic subunit of human protein kinase CK2 structurally deviates from its maize homologue in complex with the nucleotide competitive inhibitor emodin. *J Mol Biol* 377:1–8
- Raaf J, Brunstein E, Issinger OG, Niefind K (2008) The CK2 α /CK2 β interface of human protein kinase CK2 harbors a binding pocket for small molecules. *Chem. Biol.* 15:111–117
- Raaf J, Issinger O-G, Niefind K (2009) First inactive conformation of CK2 α , the catalytic subunit of protein kinase CK2. *J Mol Biol* 386:1212–1221
- Zheng J, Trafny EA, Knighton DR, Xuong NH, Taylor SS, Ten Eyck LF, Sowadski JM (1993) 2.2 Å refined crystal structure of the catalytic subunit of cAMP-dependent protein kinase complexed with MnATP and a peptide inhibitor. *Acta Crystallogr D* 49:362–365
- Bretner M, Najda-Bernatowicz A, Łebska M, Muszyńska G, Kilanowicz A, Sapota A (2008) New inhibitors of protein kinase CK2, analogues of benzimidazole and benzotriazole. *Mol Cell Biochem* 316:87–89
- Ermakova I, Boldyreff B, Issinger OG, Niefind K (2003) Crystal structure of a C-terminal deletion mutant of human protein kinase CK2 catalytic subunit. *J Mol Biol* 330:925–934
- Kabsch W (2010) XDS. *Acta Crystallogr D* 66:125–132
- McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ (2007) Phaser crystallographic software. *J Appl Crystallogr* 40:658–674
- Murshudov GN, Vagin AA, Dodson EJ (1997) Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr D* 53:240–255
- Collaborative Computational Project Number 4 (1994) The CCP4 suite: programs for protein crystallography. *Acta Crystallogr D* 50:760–763

27. Schuettelkopf AW, van Aalten DMF (2004) PRODRG: a tool for high-throughput crystallography of protein-ligand complexes. *Acta Crystallogr D* 60:1355–1363
28. Emsley P, Lohkamp B, Scott WG, Cowtan K (2010) Features and development of Coot. *Acta Crystallogr D* 66:486–501
29. Pechkova E, Zanotti G, Nicolini C (2003) Three-dimensional atomic structure of a catalytic subunit mutant of human protein kinase CK2. *Acta Crystallogr D* 59:2133–2139
30. Battistutta R, Mazzorana M, Cendron L, Bortolato A, Sarno S, Kazimierczuk Z, Zanotti G, Moro S, Pinna LA (2007) The ATP-binding site of protein kinase CK2 holds a positive electrostatic area and conserved water molecules. *Chembiochem* 8:1804–1809
31. Niefind K, Issinger OG (2010) Conformational plasticity of the catalytic subunit of protein kinase CK2 and its consequences for regulation and drug design. *Biochim Biophys Acta* 1804:484–492
32. Niefind K, Raaf J, Issinger O-G (2009) Protein kinase CK2 in health and disease: protein kinase CK2: from structures to insights. *Cell Mol Life Sci* 66:1800–1816
33. Hanks SK, Hunter T (1995) Protein kinases 6. The eukaryotic protein kinase superfamily: kinase (catalytic) domain structure and classification. *FASEB J* 9:576–596
34. Masterson LR, Cheng C, Yu T, Tonelli M, Kornev A, Taylor SS, Veglia G (2010) Dynamics connect substrate recognition to catalysis in protein kinase A. *Nat Chem Biol* 6:821–828
35. Nakaniwa T, Kinoshita T, Sekiguchi Y, Tada T, Nakanishi I, Kitaura K, Suzuki Y, Ohno H, Hirasawa A, Tsujimoto G (2009) Structure of human protein kinase CK2 α 2 with a potent indazole-derivative inhibitor. *Acta Crystallogr F* 65:75–79
36. Raaf J, Bischoff N, Klopffleisch K, Brunstein E, Olsen BB, Vilck G, Litchfield DW, Issinger OG, Niefind K (2011) Interaction between CK2 α and CK2 β , the subunits of protein kinase CK2: thermodynamic contributions of key residues on the CK2 α surface. *Biochemistry* 50:512–522
37. St-Denis NA, Derksen DR, Litchfield DW (2009) Evidence for regulation of mitotic progression through temporal phosphorylation and dephosphorylation of CK2 α . *Mol Cell Biol* 29:2068–2081
38. Esnouf RM (1997) An extensively modified version of MolScript that include greatly enhanced coloring capabilities. *J Mol Graph* 15:132–134
39. Merrit EA, Bacon DJ (1997) Raster3D: photorealistic molecular graphics. *Methods Enzymol* 277:505–524